


***Bacillus subtilis***  
**Food Grade Expression System**



**Mo Bi Tec**  
MOLECULAR BIOTECHNOLOGY



## Contents

Introduction .....	3
The Food Grade Selection System.....	3
Storage and Handling Instructions.....	5
Protocols .....	5
Media and Solutions.....	6
References.....	8
Vector Maps .....	9
Quality Warranty.....	11
Order Information, Shipping, and Storage .....	11
Contact and Support .....	11



## Introduction

Gram-positive bacteria are well known for their contributions to agricultural, medical, and food biotechnology, and for the production of recombinant proteins. Among them, *Bacillus subtilis* has been developed as an attractive host because of several reasons:

- It is non-pathogenic and is considered as a GRAS organism (generally regarded as safe), which means it can be used for food production;
- It has no significant bias in codon usage;
- It is capable of secreting functional extracellular proteins directly into the culture medium (at present, about 60 % of the commercially available enzymes are produced by *Bacillus* species);
- A large body of information concerning transcription, translation, protein folding and secretion mechanisms, genetic manipulation, and large-scale fermentation has been acquired, and the complete genome sequence of *B. subtilis* is available.

To increase the yield of heterologous proteins and to simplify operational processes, a wide variety of highly sophisticated plasmid-based expression systems have been developed. The here described **Bacillus Food Grade Expression System** was created to make the advantages of a *Bacillus* expression system also accessible to areas where antibiotic resistance gene markers are prohibited (e.g., food and feed industry). The **Bacillus Food Grade Expression System** enables stable vector-based large scale heterologous protein production by an alternative selection, without antibiotics.

The **Bacillus Food Grade Selection System** provides the following features:

- Stable high- or low-level expression without addition of any antibiotics
- All DNA contained in the final expression system is derived from *B. subtilis*
- No endotoxins are produced
- No inclusion bodies are formed
- Protease-deficient strain for producing secretory enzymes is available

## The Food Grade Selection System

The **Bacillus Food Grade Selection System** is based on the interplay of an endogenous *Bacillus* toxin **EndoA** and its **antitoxin EndoB**.

EndoA (encoded by *ydcE*) is an endoribonuclease that specifically cleaves mRNA at a five Base U↓ACAU sequence (Pellegrini *et al.*, 2005, Park *et al.*, 2011). During normal growth conditions EndoA is inactivated by forming a heterohexameric complex with its cognate antitoxin EndoB (encoded by *ydcD*; Simanshu *et al.*, 2013). Since the antitoxin is relatively unstable, it is essential for the cell to continuously produce sufficient amounts of EndoB to inactivate the more stable toxin. These characteristics are utilized for the here described selection system.

### The mechanism of the selection system

The antitoxin encoding gene *ydcD* is located within the expression vector under the control of a constitutive *Bacillus* promoter whereas the toxin expression cassette is integrated into the genome and is controlled by a xylose-inducible promoter. If the expression of the toxin is induced by xylose but the plasmid with the encoded antitoxin gets lost, no more antitoxin



can be produced, leading to an active endoribonuclease that results in inhibition of protein synthesis, cell growth arrest, and finally dying of the cell.

This expression system has been proven to be very stable, with no plasmid loss observed in 100 generations. Compared with traditional antibiotic-dependent expression systems, this system results in greater biomass and higher titers of the desired products. This has been shown for the expression of the green fluorescent protein and the metabolic product hyaluronan, respectively (Yang *et al.*, 2016).

### The expression vectors

Dependent on the particular requirements there can be chosen between two *B. subtilis* expression vectors **pTTB1** and **pTTB2**. These vectors differ in their origin of replication and copy number concerning *B. subtilis*. **pTTB1** is a low copy number vector, replicating via theta replication modus, whereas **pTTB2** is high copy number, replicating by rolling circle mechanism.

Both vectors share the following features:

- For easier handling the vectors are designed as *B. subtilis* / *E. coli* shuttle vectors. The parts of the vector used for cloning with *E. coli* (*E. coli* origin ColE1 derivative and ampicillin resistance cassette Amp) can be eliminated afterwards by restriction enzyme cleavage and religation of the vector (for details see below). This technique connects the advantage of easy cloning (with *E. coli*) with the food grade property of *B. subtilis*.
- For food grade selection with *B. subtilis* the antitoxin-encoding gene *ydcD* is included under control of a constitutive *Bacillus* promoter (the strong  $P_{ylyxM}$  promoter in pTTB1 and the weaker  $P_{aadD}$  promoter in pTTB2).
- A multiple cloning site for cloning the gene of interest downstream of the constitutive promoter  $P_{43}$  is provided.

### The food grade *Bacillus* strains

For expressing the gene of interest under food grade conditions, two *B. subtilis* strains are available; TEA and WEA. Both strains contain the toxin expression cassette *ydcE* under control of the xylose-inducible promoter  $P_{xyl}$ , whereas the former *ydcDE* operon (containing the toxin and antitoxin gene) is deleted. The strain *B. subtilis* TEA is based on *B. subtilis* 168 Marburg and is recommended for intracellular protein expression and pathway engineering. *B. subtilis* WEA originated from the eightfold extracellular protease-deficient strain *B. subtilis* WB800N and is particularly constructed for secretory protein production. Both strains are suitable hosts for both the food grade low copy expression vector pTTB1 and the high copy expression vector pTTB2.

**TEA:** *trpC2 ydcDE::P<sub>xyl</sub>-ydcE*

**WEA:** *nprE aprE epr bpr mpr::ble nprB::bsrΔvpr wprA::hyg cm::neo ydcDE::P<sub>xyl</sub>-ydcE; NeoR*

Please note: this strain carries resistance to neomycin!



## Storage and Handling Instructions

### Storage and handling of plasmids

Plasmids are supplied lyophilized. Upon receipt, add 100 µl distilled water (final concentration 0.1 µg/µl) and incubate at 50 °C for 5 minutes. Vortex for 1 minute and store at -20 °C. Please note that all plasmids of this system are *E. coli* / *B. subtilis* shuttle vectors.

### Storage and handling of *Bacillus* strains

The *Bacillus* strains are supplied as frozen cultures and shipped on dry ice. Store the stock at -80 °C. For propagation remove tube from freezer, scratch off some material from the surface of the frozen stock using a sterile loop. Streak onto an LB plate (see page 7), seal the plate with parafilm, and incubate at 37 °C overnight. *Bacillus* plates can be stored at 4 °C for 1 month. Use fresh bacteria, starting from one single colony for preparing competent cells.

## Protocols

### Growth conditions

Detailed protocols for *E. coli* and *B. subtilis* molecular genetic handling (growth, transformation, etc.) can be found in the relevant laboratory manuals such as Sambrook and Russell (2001).

*B. subtilis* and *E. coli* can be grown aerobically at 37 °C in 2xYT medium (Bagyan *et al.*, 1998). Under optimal conditions the doubling time of *E. coli* is 20 min, of *B. subtilis* 30 min. For selecting transformed *E. coli* cells, use 50-100 µg/ml ampicillin. For selecting transformed *Bacillus* TEA and WEA strains, add xylose to a final concentration of 2 g/L to the medium.

### Vector propagation and cloning the DNA fragment of interest

Since both expression vectors (pTTB1 and pTTB2) are designed as *E. coli* / *B. subtilis* shuttle vectors, we recommend using *E. coli* for plasmid propagation and for cloning the DNA fragment of interest. Follow standard protocols for propagation in *E. coli*, *E. coli* plasmid mini preparation, restriction endonuclease cleavages, ligation of the desired DNA fragment into the vector, and transformation of *E. coli* (Sambrook and Russell, 2001). For selecting successfully transformed *E. coli* cells use 50-100 µg ampicillin. To retain food grade properties both vectors provide the opportunity to remove the functional *E. coli* parts (i.e., ampicillin resistance cassette and origin for replication) after finishing the construct. These parts can be removed by cleavage with one single restriction enzyme (for pTTB1 use *SpeI*, for pTTB2 use *EcoRI*) followed by purification of the larger fragment and religation. The religated vector does now only contain DNA that is derived exclusively from *B. subtilis* (with the exception of the cloned fragment).





- 2xYT medium: 16 g tryptone  
10 g yeast extract  
5 g sodium chloride (NaCl)  
add distilled water to 1000 ml and autoclave  
(121 °C, 15 min)
- 2xYT / xylose agar plates: add 1.5 % agar to the 2YT medium before  
autoclaving, let cool down to 70 °C and add 2 %  
sterile-filtered xylose before pouring the plates
- 10x S-base (Spizizen's salt): 2 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
14 g K<sub>2</sub>HPO<sub>4</sub>  
6 g KH<sub>2</sub>PO<sub>4</sub>  
1 g sodium citrate  
add distilled water to 100 ml and autoclave  
add 0.1 ml 1 M MgSO<sub>4</sub> after autoclaving
- HS medium: 66.5 ml distilled water  
10.0 ml 10x S-base  
2.5 ml 20 % (w/v) glucose  
5.0 ml 0.1 % (w/v) L-tryptophan  
1.0 ml 2 % (w/v) casamino acids  
5.0 ml 10 % (w/v) yeast extract (Difco)  
10.0 ml 8 % (w/v) arginine, 0.4 % histidine  
autoclave all components separately  
tryptophan solution: sterile filtration
- LS medium: 80.0 ml distilled water  
10.0 ml 10x S-base  
2.5 ml 20 % (w/v) glucose  
0.5 ml 0.1 % (w/v) L-tryptophan  
0.5 ml 2 % (w/v) casamino acids  
5.0 ml 2 % (w/v) yeast extract (Difco)  
0.25 ml 1 mM MgCl<sub>2</sub>  
0.05 ml 1 mM CaCl<sub>2</sub>  
autoclave all components separately  
tryptophan solution: sterile filtration
- 0.1 M EGTA dissolve 3.8 g EGTA in 50 ml distilled water  
adjust the pH to 7.2 using 10 N NaOH  
add distilled water to 100 ml; autoclave




## References

- Bagyan I. *et al.* (1998). The *katX* Gene, which codes for the catalase in spores of *Bacillus subtilis*, is a forespore-specific gene controlled by sigmaF, and KatX is essential for hydrogen peroxide resistance of the germinating spore; *J Bacteriol.* 180(8), 2057–2062
- Klein C. *et al.* (1992). Analysis of genes involved in biosynthesis of the lantibiotic subtilin; *Appl. and Environ. Microbiol.* 58, 132-142
- Park *et al.* (2001). *Bacillus subtilis* MazF-bs (EndoA) is a UACAU-specific mRNA interferase; *FEBS Lett.* 585, 2526-2532
- Pellegrini *et al.* (2005). The *Bacillus subtilis* ydcDE operon encodes an endoribonuclease of the MazF/Penk family and its inhibitor; *Mol Microbiol.* 56, 1139-1148
- Sambrook, J. & Russell, D.W. (2001). *Molecular cloning: a laboratory manual.* (New York: Cold Spring Harbor Laboratory Press).
- Simanshu D.K. *et al.* (2013). Structural basis of mRNA recognition and cleavage by toxin MazF and its regulation by antitoxin MazE in *Bacillus subtilis*; *Mol.Cell* 52, 447-458
- Yang S. *et al.* (2016). Construction of a novel, stable, food-grade expression system by engineering the endogenous toxin-antitoxin system in *Bacillus subtilis*; *J. Biotechnol.* 219, 40-47



## Vector Maps

### Vector map pTTB1 (low copy)




	Type	Start	End	Name	Description
	Selectable genetic marker	724	443	ydcD	Antitoxin encoding gene
	Promoter	1054	755	PylxM	Strong constitutive promoter for expression of ydcD
	Promoter	1067	1365	P43	Constitutive promoter for expression of gene of interest
	MCS	1366	1395	MCS	Multiple Cloning Site
	Gene	2388	3422	repA	Gene of replication protein RepA ( <i>B. subtilis</i> )
	Origin of replication	4918	5517	ColE1 ori*	origin of replication ( <i>E. coli</i> ); ColE1 incompatibility group
	Selectable genetic marker	6539	5679	Amp	Ampicillin resistance ( <i>E. coli</i> )

The complete DNA sequence is available at our webpage [www.mobitec.com](http://www.mobitec.com).



## Vector map pTTB2 (high copy)



	Type	Start	End	Name	Description
	Gene	814	1818	repU	Gene of replication protein RepU ( <i>B. subtilis</i> )
	Promoter	1054	755	PaadD	constitutive promoter for expression of ydcD
	Selectable genetic marker	1987	2268	ydcD	Antitoxin encoding gene
	Promoter	3674	3947	P43	constitutive promoter for expression of gene of interest
	MCS	3948	3990	MCS	Multiple Cloning Site
	Origin of replication	4409	5008	ColE1 ori*	origin of replication ( <i>E. coli</i> ); ColE1 incompatibility group
	Selectable genetic marker	6030	5170	Amp	Ampicillin resistance ( <i>E. coli</i> )

Please consider: EcoRI is no single cutter!

The complete DNA sequence is available at our webpage [www.mobitec.com](http://www.mobitec.com).



## Quality Warranty

The vector features and restriction sites specified in this manual are verified by sequencing or checked for being functional. MoBiTec does not give any guarantee for sequence data of nonfunctional parts of the vectors.

## Order Information, Shipping, and Storage

Order#	Product	Amount
PBS041	pTTB1	10 µg
PBS042	pTTB2	10 µg
Shipped at RT; plasmids are lyophilized from water and can be stored at 4 °C. Once the DNA has been dissolved in sterile buffer or water, store at -20 °C.		
PBS043	Bacillus subtilis strain TEA	1 ml
PBS044	Bacillus subtilis strain WEA	1 ml
Shipped on dry ice; store at -20 °C.		

The *Bacillus subtilis* Food Grade Expression System was developed by researchers of the Jiangnan University, China (Yang *et al.*, 2016). All strains and vectors belonging to this system are for academic research and development only.

Commercial enterprises: To obtain strains and plasmids of this system, a license needs to be negotiated. For license agreements please mail to [info@mobitec.com](mailto:info@mobitec.com).

## Contact and Support

MoBiTec GmbH ♦ Lotzestrasse 22a ♦ D-37083 Goettingen ♦ Germany

**Customer Service** – General inquiries & orders  
 phone: +49 (0)551 707 22 0  
 fax: +49 (0)551 707 22 22  
 e-mail: [order@mobitec.com](mailto:order@mobitec.com)

**Technical Service** – Product information  
 phone: +49 (0)551 707 22 70  
 fax: +49 (0)551 707 22 77  
 e-mail: [info@mobitec.com](mailto:info@mobitec.com)

**MoBiTec in your area:** Find your local distributor at [www.mobitec.com](http://www.mobitec.com)